

Pars Quarta. 115

tionem subministrarem: & quia incertum est, correxerintne Arabes in Ptolemaicis istis aliqua, an omnium veriorum diversitas à transcriptorum incuria sit orta. Pauca sunt, quibus ego manum admovi, vel in libro de stella Serpentarij vel alijs, quas caractere antiquo, monendi causa, expressi.

DENOMINATIO STELLARUM.		Longitudo G. M. S.	Latitudo G. M.	Magnitudo	DENOMINATIO STELLARUM.		Longitudo G. M. S.	Latitudo G. M.	Magnitudo	
In URS A minore informis Clavij (signo falso) est 8 Tychoens.					Proximè infra ultimam	10.22	25.2	B	6	
In URS A majore 39. 40. 41. in meo Exemplari sunt magni.					Sequens manum Ophiuchi	29.45	16.0		5	
non 3, sed 5, lat. 34. 59. 21. 38, 20. 44, & Ptol. 8 uam.					In ANDROMEDA In Syriate	4.0	34.40		5	
formaturum inter eas esse videtur, quia Trapez. unius pro o habet o.					Austrina [duarum bor.]	5.30	32.40		5	
In CEPHEO, informis pr. Tiarum	4.30	V	64.0	B	5	3.0	V	44.0	B	3
In BOOTE, 10 Ptol. in venabulo	28.45	II	45.45	4	In ARIETE. Trium iusfor. borea	12.20	12.44		B	5
In ENGONASI in coxa sinist. pr.	1.20	II	55.55	5	In extremo pede posteriori	6.0	8	5.10	A	4
Sequens	2.30	II	58.15	5	In TAURO. Observatio de Plejade.					
Informis dextri brachij Australior	24.39	M	57.30	5	Commendat eam creber appulsus Lunæ, & usus in					
In OLORE in dextro genu	3.0	M	63.20	5	obs. antiquissima magni momenti. Igitur Tycho solas 4					
In PERSEO, informis à sinistro genu ad ortum	2.10	II	17.50	B	ne. Enumerat: & tamen primam earum non invenio in Observationibus. Etiamq; meam Msc. pro 23. 13. 15, habet					
In OPHIUCHO & desunt multæ Ptolemaicæ, & suspectæ sunt aliqua, quas Clavius ex Pifero sumpsit, qui alijs merum Catalogum Tychoenicum complexus est: cum tamen hæc non ferat tantam. Idem Ptol. exortum tamen desinit in meo exemplari, Quarum altitudines & culminationes cum instrumentis ceperim; Catalogum æquiculus à quadrilatero, deficit in ejus enumeratione, quartò ad ex libro meo de stella nova transcribam, nihil præjudicans. cio exprimit. At meæ dispositiones diversissimorum temporum, tribus Tychoenicis innixæ, consentiunt satis propinquè in hos numeros; ac si sic in Ptol. sit legendum:										
In recta trium ante pedè præ. bor.	25.42	M	9.27	B	5	24. 5	8	4.2	I	
Media	24.47	M	6.42	5	5	23.50	8	4.13		
Infima	24. 5	M	3.45	5	6	24.24	8	4. 3		
Quæ præcedit genu anterius	1. 3	II	13.34	5	4	24. 4	II	3.55		
Duarum inter pedes superior	6.51	II	11.50	6	4	24. 7	8	4.25		
Inferior	5.49	II	4.38	6	5	24. 0	8	4.26		
Trium in crure anteriori borea	2.47	II	5.42	4	4	24.47	8	3.55		
o. Media	2. 7	II	3.11	4	4					
p. Australissima	1.45	II	1.38	4	4					
q. In calcaneo pedis hujus	4. 4		0.26	4	4					
In femore sequenti	15.39		17.28	5	4					
m. In cubito sequenti	19. 2	II	14.57	4	4					
n a. In pedis sequentis Tibia	15.42	II	1.57	B	4					
a. In digito pedis	14. 7	II	3.37	A	4					
In dorso pedis	14.47	II	1.27	5	4					
b. In Plantâ, clara	15.47	II	1.43	3	4					
c. In vola	16.37	II	0.59	4	4					
d. e. In calce seu talo	18. 5	II	0.57	A	4					
Sequens pedem hunc	20. 4	II	1.22	B	4					
Infor. inter hum. Oph. & Herc.	26.27	M	35. 0	B	4					
b b. Ad humerum Oph. in Rhomborealissima	25. 7	II	28. 0	4	4					
c c. Mediarum præcedens	24.57	II	26.40	4	4					
d d. Sequens	26.50	II	26.28	4	4					
e e. Infima	25.40	II	24.45	4	4					
Stella nova anni 1604	17.40	II	1.56	4	4					
Supra informatas solitaria	27.50	II	32.45	B	4					
In SERPENTE. Quæ supra duplicem, octava Ptolemaeo	16.33	M	26.36	B	4					
In crista	18. 9	M	37.15	5	4					
In spira, apex Isoscelis 13 Ptol.	15. 0	II	10.25	4	4					
Basis præcedens 14 Ptol.	19.23	II	8. 4	4	4					
Sequens 15 Ptol.	20. 9	II	10.23	4	4					
Supra caudam duarum superior	29.49	II	26.38	6	4					
Inferior	1.39	II	23.28	6	4					
Infra caudam clara	3. 0	II	15.49	4	4					
Trium inter ultimas caudæ ima	2. 0	II	21.29	6	4					
Medis	4. 6	II	22.42	6	4					
Postrema	6.40	II	24.52	B	6					

270

268

274

264

259

219

169

Ende

Anfang